

Orbits: inhomogeneous fields and fluid currents

Alfio Bonanno

INAF – Osservatorio Astrofisico di Catania

Outline

1. Inhomogeneous fields
2. From particles to fluid
3. Validity of the guiding-centre picture

Reading

Chen, ch. 2

Sturrock, ch. 3

Goldston & Rutherford, ch. 2–3

Northrop (1963)

Notes: chapter 3, §3.3–3.5

Problems: set 2

Where we are

Key result: recap

- ▶ gyration at ω_c on a circle r_L around the guiding centre
- ▶ magnetic moment $\mu = \frac{1}{2}mv_{\perp}^2/B$, antiparallel to $\mathbf{B}$
- ▶ any force $\mathbf{F}$: drift $(c/q)\mathbf{F} \times \mathbf{B}/B^2$; $\mathbf{E} \times \mathbf{B}$ for all species
- ▶ today: fields that vary in space, and how drifts add up to fluid currents

Inhomogeneous fields

The grad- B drift (1): orbit average

$\mathbf{B} = B_z(x) \hat{\mathbf{z}}$, with $r_L |\nabla B| \ll B$. Expand about the guiding centre x_0 :

$$F_x = \frac{q}{c} v_y B_z(x) = \frac{q}{c} v_y B_0 + \frac{q}{c} v_y (x - x_0) \frac{\partial B}{\partial x}$$

Average over the unperturbed orbit: $v_x = v_\perp \cos \Omega t$, $v_y = -v_\perp \sin \Omega t$,
 $x - x_0 = (v_\perp / \Omega) \sin \Omega t$:

$$\langle v_y \rangle = 0, \quad \langle v_y (x - x_0) \rangle = -\frac{v_\perp^2}{\Omega} \langle \sin^2 \Omega t \rangle = -\frac{v_\perp^2}{2\Omega} \Rightarrow \langle F_x \rangle = -\frac{q}{c} \frac{v_\perp^2}{2\Omega} \frac{\partial B}{\partial x} = -\frac{\frac{1}{2} m v_\perp^2}{B} \frac{\partial B}{\partial x}$$

Key result

$$\langle \mathbf{F} \rangle = -\mu \nabla B \quad (\langle F_y \rangle = 0; \text{ a diamagnetic dipole, pushed to weak field})$$

The grad- B drift (2)

Insert $\langle \mathbf{F} \rangle = -\mu \nabla B$ in the force drift:


$$\mathbf{v}_{\nabla B} = \frac{c}{q} \frac{(-\mu \nabla B) \times \mathbf{B}}{B^2} = \frac{\mu c}{q} \frac{\mathbf{B} \times \nabla B}{B^2}$$

Key result: grad- B drift

$$\mathbf{v}_{\nabla B} = \frac{v_{\perp}^2}{2\Omega} \frac{\hat{\mathbf{b}} \times \nabla B}{B}, \quad v_{\nabla B} \sim \frac{v_{\perp}}{2} \frac{r_L}{L_B}$$

∇B drift: opposite for $\pm q$

- ▶ smaller than $v_{\perp}$ by r_L/L_B
- ▶ opposite for $\pm q$: **current**
- ▶ radius smaller where B is larger: orbit does not close



The curvature drift

Particle sliding along a curved line at $v_{\parallel}$: centrifugal force in its frame,

$$\mathbf{F}_{\text{cf}} = \frac{mv_{\parallel}^2}{R_c} \hat{\mathbf{e}}_r = mv_{\parallel}^2 \frac{\mathbf{R}_c}{R_c^2}$$

$\mathbf{R}_c$: from the centre of curvature to the line.

Key result: curvature drift

$$\mathbf{v}_R = \frac{c}{q} \frac{\mathbf{F}_{\text{cf}} \times \mathbf{B}}{B^2} = \frac{mcv_{\parallel}^2}{q} \frac{\mathbf{R}_c \times \mathbf{B}}{R_c^2 B^2} = \frac{v_{\parallel}^2}{\Omega} \hat{\mathbf{b}} \times \boldsymbol{\kappa}, \quad \boldsymbol{\kappa} = (\hat{\mathbf{b}} \cdot \nabla) \hat{\mathbf{b}} = -\frac{\mathbf{R}_c}{R_c^2}$$

A curved field always has a gradient too. Next: both together in a vacuum field.

Curvature + grad- B in a vacuum field

Vacuum or low pressure: $\nabla \times \mathbf{B} = 0$. Local cylindrical coordinates, $\mathbf{B} = B_\theta(r)\hat{\theta}$:

$$(\nabla \times \mathbf{B})_z = \frac{1}{r} \frac{d}{dr}(rB_\theta) = 0 \Rightarrow B_\theta \propto \frac{1}{r} \Rightarrow \frac{\nabla B}{B} = -\frac{\hat{\mathbf{e}}_r}{r} = -\frac{\mathbf{R}_c}{R_c^2}$$

Then $\mathbf{v}_{\nabla B} = \frac{mc v_\perp^2}{2qB} \frac{\mathbf{B} \times \nabla B}{B^2} = \frac{mc v_\perp^2}{2q} \frac{\mathbf{R}_c \times \mathbf{B}}{R_c^2 B^2}$: same direction as $\mathbf{v}_R$.


Key result

$$\mathbf{v}_R + \mathbf{v}_{\nabla B} = \frac{mc}{q} \frac{\mathbf{R}_c \times \mathbf{B}}{R_c^2 B^2} \left(v_\parallel^2 + \frac{1}{2} v_\perp^2 \right)$$

Maxwellian: $\langle v_\parallel^2 + \frac{1}{2} v_\perp^2 \rangle = T/m + T/m = 2T/m$, so $\langle \mathbf{v}_R + \mathbf{v}_{\nabla B} \rangle = \frac{2cT}{qBR_c} \propto \frac{T}{BR_c}$,

independent of mass.

Why a simple torus cannot confine a plasma



Field lines are circles around the axis ($B \propto 1/R$):

1. curvature + grad- B : ions up, electrons down
2. charge separation $\Rightarrow$ vertical E
3. $v_E = cE \times B/B^2$ **outwards**, for both species
4. the plasma hits the wall

Estimate

$T = 1 \text{ keV}$, $B = 10^4 \text{ G}$, $R_c = 1 \text{ m}$: $v = 2cT/eBR_c \approx 2 \text{ km s}^{-1}$: the wall is reached in $\sim 1 \text{ ms}$.

Cure: twist the field lines into helices, so particles spend equal times above and below the midplane. Tokamak (plasma current), stellarator (external coils).

From particles to fluid

A paradox

Uniform $\mathbf{B} = B\hat{z}$, density gradient along x , uniform T .

Particle view

uniform B : no grad- B , no curvature drift.

No guiding centre moves.

Fluid view


force balance $\mathbf{j} \times \mathbf{B}/c = \nabla p$ requires

$$\mathbf{j} = c \frac{\hat{\mathbf{b}} \times \nabla p}{B} \neq 0$$

Key result

Where does the current come from? From the **gyration** itself: each particle is a current loop, and loops do not cancel where their density varies.

The magnetisation current



Across a vertical line: loops on the left contribute their (downward) right sides, loops on the right their (upward) left sides. More loops on the right: net upward current, along $\hat{\mathbf{b}} \times \nabla n$.

Magnetisation = moment per unit volume, with $p_{\perp} = \sum n \langle \frac{1}{2} m_s v_{\perp}^2 \rangle$:

$$\mathbf{M} = \sum_s n_s \langle \boldsymbol{\mu}_s \rangle = - \sum_s n_s \left\langle \frac{\frac{1}{2} m_s v_{\perp}^2}{B} \right\rangle \hat{\mathbf{b}} = - \frac{p_{\perp}}{B} \hat{\mathbf{b}}, \quad \mathbf{j}_M = c \nabla \times \mathbf{M}$$

Summing all contributions

Straight lines, $\mathbf{B} = B(x)\hat{\mathbf{z}}$, isotropic $\rho(x)$. **Drift current** (grad- B , $\mathbf{v}_{\nabla B} = (\mu c/q) \hat{\mathbf{b}} \times \nabla B/B$):

$$\mathbf{j}_{\nabla B} = \sum nq \mathbf{v}_{\nabla B} = c \left(\sum n\mu \right) \frac{\hat{\mathbf{b}} \times \nabla B}{B} = c\rho \frac{\hat{\mathbf{b}} \times \nabla B}{B^2}$$

Magnetisation current, $\mathbf{M} = -(p/B)\hat{\mathbf{z}}$, $\nabla \times (f\hat{\mathbf{z}}) = \nabla f \times \hat{\mathbf{z}}$:

$$\mathbf{j}_M = -c \nabla \left(\frac{p}{B} \right) \times \hat{\mathbf{z}} = c \frac{\hat{\mathbf{b}} \times \nabla p}{B} - c\rho \frac{\hat{\mathbf{b}} \times \nabla B}{B^2}$$

Key result: diamagnetic current

$$\mathbf{j}_{\perp} = \mathbf{j}_{\nabla B} + \mathbf{j}_M = c \frac{\hat{\mathbf{b}} \times \nabla p}{B}, \quad \frac{\mathbf{j}_{\perp} \times \mathbf{B}}{c} = (\hat{\mathbf{b}} \times \nabla p) \times \hat{\mathbf{b}} = \nabla_{\perp} p$$

Particles and fluid agree

- ▶ the grad- B terms cancel exactly: the total current depends only on ∇p
- ▶ particle and fluid pictures agree **only** when the magnetisation current is included
- ▶ the fluid velocity is *not* the guiding-centre velocity

Key result

Two-fluid language: each species carries the **diamagnetic drift**

$$\mathbf{u}_{*s} = c \frac{\hat{\mathbf{b}} \times \nabla p_s}{q_s n_s B}$$

a fluid velocity that no individual guiding centre has (ch. 9).

Curved fields add $c(p_{\parallel} - p_{\perp}) \hat{\mathbf{b}} \times \boldsymbol{\kappa} / B$, zero for isotropic p .

Validity of the guiding-centre picture

Ordering and summary of drifts

Small parameters: $\epsilon = r_L/L \ll 1$, $\omega/\omega_c \ll 1$. Drifts are first order:
 $v_{\nabla B}/v_{\perp} \sim v_R/v_{\parallel} \sim \epsilon$, $v_P/v_E \sim \omega/\omega_c$.

drift	velocity	depends on q ?	current?
$\mathbf{E} \times \mathbf{B}$	$c \mathbf{E} \times \mathbf{B}/B^2$	no	no
force $\mathbf{F}$	$(c/q) \mathbf{F} \times \mathbf{B}/B^2$	yes	yes
gravitational	$(mc/q) \mathbf{g} \times \mathbf{B}/B^2$	yes, $\propto m$	yes (ions)
grad- B	$(v_{\perp}^2/2\Omega) \hat{\mathbf{b}} \times \nabla B/B$	yes	yes
curvature	$(v_{\parallel}^2/\Omega) \hat{\mathbf{b}} \times \boldsymbol{\kappa}$	yes	yes
polarisation	$(1/\Omega)(c/B) \partial_t \mathbf{E}_{\perp}$	yes, $\propto m$	yes (ions)

Summary

Guiding-centre velocity to first order:

$$\mathbf{v}_{gc} = v_{\parallel} \hat{\mathbf{b}} + \mathbf{v}_E + \frac{\hat{\mathbf{b}}}{\Omega} \times \left(\frac{v_{\perp}^2}{2} \frac{\nabla B}{B} + v_{\parallel}^2 \boldsymbol{\kappa} + \frac{d\mathbf{v}_E}{dt} \right)$$

- ▶ gyration: $\omega_c = |q|B/mc$, $r_L = v_{\perp}/\omega_c$,
 $\mu = \frac{1}{2} m v_{\perp}^2 / B$
- ▶ $\mathbf{E} \times \mathbf{B}$: no current; all other drifts carry currents
- ▶ drift + magnetisation currents = $c \hat{\mathbf{b}} \times \nabla p / B$

Still missing: the motion *along* $\mathbf{B}$, governed by the mirror force $-\mu \partial B / \partial s$ and the conservation of μ .

Next: module 4a, magnetic mirrors; then radiation belts and Fermi acceleration.

Problems: set 2.